

Alexandria Location

4660 Kenmore Ave, Suite 1120
Alexandria, VA 22304

tel: 703-680-2111
fax: 703-878-3939

Woodbridge Location

2296 Opitz Blvd, Suite 350
Woodbridge, VA 22191

Alok Desai, M.D. Pratik Desai, M.D. Nilay Gandhi, M.D. John Klein M.D. Inderjit Singh M.D.

Ureteral Stent Instructions

WHAT IS A STENT?

- Soft, hollow tube with multiple holes
- Allows temporary drainage if kidney is blocked
- It is not visible from the outside of the body
- Remains for 1-2 weeks or long-term for several months

PROCEDURE

- No cuts/incisions will be made on your skin
- A camera is inserted into the urethra and bladder
- The stent is inserted and an x-ray confirms the position

ACTIVITY

- Resume your normal routine after the stent is inserted
- Rest if you notice pain or blood in the urine with activity
- It is ok to engage in sexual activity

INSTRUCTIONS

- Drink 64 ounces of fluids a day
- If prescribed, take the pain medication, spasm medication, and/or antibiotic as directed.

EXPECTED SYMPTOMS

- Urge to urinate
- Urinary frequency
- Urinary burning/dysuria
- Blood in urine/hematuria
- Pain when/after urinating (back, side, groin)

WHEN TO CALL THE DOC

- Fever > 100.5° F and/or chills
- Nausea/vomiting
- Cloudy urine, blood clots
- Unable to urinate, urinary leakage
- Pain not controlled with medication

FOLLOW-UP CARE

- The ureteral stent is **NOT PERMANENT**. It must be removed/changed so you don't get seriously ill or permanently damage your kidney
- Please discuss the plan for your stent with your urologist **within 1-2 weeks** of placement

